

		REPORT (DOCUMENTATI	1 1 2 2 2 4 1 4 2 1 2 1 2 2 2 2 2 2 2 2			n Approved 8 No. 0704-0188		
1a REPORT S	ECURITY CLASS	SIFICATION							
	ssified			<u> </u>					
2a SECURITY	CLASSIFICATIO	N AUTHORITY		3 DISTRIBUTION Approved	for public		and sale:		
26 DECLASSIFICATION DOWNGRADING SCHOOLES				distribution unlimited.					
4 PERFORMIN	IG ORGANIZAT	ION REPO NUMBE	MST	5 MONITORING	ORGANIZATION RE	PORT NUMBE	R(S)		
Repor	t #21		8	4135010					
6a NAME OF	PERFORMING	ORGANIZATION	6b OFFICE SYMBOL (If applicable)	7a NAME OF MONITORING ORGANIZATION					
Unive	rsity of 1	Minnesota	ONR	Office of Naval Research					
6c. ADDRESS	(City, State, an	d ZIP Code)		7b ADDRESS (City, State, and ZIP Code)					
		Chemistry		000 % 0 4					
	Leasant Si			800 N. Quincy Street					
Minnea	apolis, M	N 55455-0431		Arlington, VA 22217-5000					
	FUNDING / SPC	ONSORING	8b OFFICE SYMBOL	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER					
ORGANIZATION (If applicable) Office of Naval Research ONR			N001489J1301						
	City, State, and		<u> </u>	10 SOURCE OF FUNDING NUMBERS					
	. Quincy S			PROGRAM ELEMENT NO	PROJECT NO	TASK NO	WORK UNIT ACCESSION NO		
		2217-5000		61153N	4135		4135010		
11 TITLE (Include Security Classification)							4131010		
"Chromatographic Assessment of Hydrogen-Bond Donating Ability" (Unclassified)									
12 PERSONAL		r M.C. Car	r P Li						
Wanke, R.; Etter, M.C.: Carr, P.W. 13a TYPE OF REPORT (Year, Month, Day) 15 PAGE COUNT									
Technical FROM 6/1/92 TO 5/31/91 April 22, 1993 8									
16 SUPPLEMENTARY NOTATION									
17	COSATI	CODES	18 SUBJECT TERMS (C	Continue on reverse	if necessary and	identify by b	lock number)		
FIELD	GROUP	SUB-GROUP	1. 1.	1 1/1					
07	03		nyarogen	bond/chromatography					
					· · · · · · · · · · · · · · · · · · ·				
19 ABSTRACT	(Continue on	reverse if necessary	and identify by block no	ımber)					
Generally, hydrogen bonding preferentially occurs between the best donors and best acceptors in the organic solid state. Knowledge of the relative hydrogen-bond acidity and basicity can, therefore, be used in molecular crystal engineering. A liquid chromatography method has been developed to assess the hydrogen-bond acidity of potential hydrogen-bond donors used in cocrystallizations. Hydrogen-bond donor solutes are chromatographed on a poly(vinylpyridine-divinylbenzene) column under two different sets of mobile phase conditions. The relative retention is proportional to the hydrogen-									
bond acidity of the solute and therefore, provides an a priori measure of the hydrogen-									
bond acidity of a potential cocrystal component.									
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT 21. ABSTRACT SECURITY CLASSIFICATION									
☐ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT ☐ DTIC USERS ☐ Unclassified 22a NAME OF RESPONSIBLE INDIVIDUAL 22b TELEPHONE (Include Area Code) 22c OFFICE SYMBOL									
				22b TELEPHONE (Include Area Code) 22c OFFICE SYMBOL ONR Code 1113					
narol	d E. Guar	. <u>u</u>		1,2021,646	-4311	1 July	~~~		

OFFICE OF NAVAL RESEARCH

Grant or Contract N001489J1301

R&T Code 4135010

Technical Report #21

"Chromatographic Assessment of Hydrogen-Bond Donating Ability

by

R. Wanke, M.C. Etter, and P.W. Carr

University of Minnesota Department of Chemistry Minneapolis, MN

April 22, 1993

Reproduction in whole or in part is permitted for any purpose of the United States Government

This document has been approved for public release and sale; its distribution is unlimited.

93-10499

93 5 11 16 6

Hydrogen bonds are strong orienting forces in the organic solid state. A general heuristic principle that has guided our cocrystallization studies is "the best hydrogen-bond donor hydrogen bonds to the best hydrogen-bond acceptor". Successive hydrogen-bond donors, barring severe conformational or crystal packing constraints, hydrogen bond to available acceptors in rank order. Therefore an a priori knowledge of the relative hydrogen-bond strength of the hydrogen-bonding groups within a given cocrystal should allow for the prediction of the hydrogen-bond connectivity pattern in the solid state.

Within our laboratory, Taft ß values² have been used to assess the hydrogen-bond basicity and pKa values have been used to estimate the hydrogen-bond acidity of compounds or specific functional groups within a cocrystal pair. No general correlation exists between pKa and the free energy of hydrogen-bond association across different families of hydrogen-bond donors; therefore, the use of pKa to assess the relative hydrogen-bond acidity of donors is prone to inaccuracy. Relative scales of hydrogen-bond acidity such as the α_2^H scale³, which have been derived from the free energies of hydrogen-bond association in carbon tetrachloride, do exist. However, tabulations of α_2^H are less extensive than ß and do not include solid compounds of prime interest in cocrystallization studies. Therefore, chromatography was selected as a potential empirical method to assess the hydrogen-bond acidity of hydrogen-bond donor compounds used in cocrystallization studies.

In chromatography, small differences in molecular interaction between a solute for the stationary and mobile phases result in macroscopic differences in chromatographic retention. The logarithm of the chromatographic capacity factor (log k') is proportional to the free energy of partitioning of the solute from the mobile phase into the stationary phase and as such, accounts for specific hydrogen-bonding interactions and van der Waal's interactions of the solute with the stationary or mobile phase. In order to isolate the hydrogen-bonding contribution to the retention, two LC approaches were devised that involve comparing the retention of a given solute on two matched stationary phases or on two matched mobile phases, respectively. The columns and mobile phases for the two different approaches are carefully selected to be matched in their nonhydrogen-

Dist Special

DITO CALLED BORGERO B

Etter, M.C. J. Phys. Chem. 1991, 95, 4601.

² Kamlet, M.J.; Abboud, J-L. M.; Abraham, M.H.; Taft, R.W. J. Org. Chem. 1983, 48, 2877.

Abraham, M.H.; Grellier, P.L.; Prior, D.V.; Duce, P.P.; Morris, J.J.; Taylor, P.J. J. Chem. Soc., Perkin Trans. 2 1989, 699.

bonding properties but different in their hydrogen-bond accepting abilities. As shown in Scheme 1, two polymeric columns, a poly(vinylpyridine-divinylbenzene) column and a poly(styrene-divinylbenzene) column, along with a 90:10 acetonitrile/dimethylsulfoxide mobile phase are implemented for the dual column approach. For the dual mobile phase approach, solutes are chromatographed on the poly(vinylpyridine-divinylbenzene) column using two binary mobile phases, a 95:5 acetonitrile/triethylphosphate and a 80:20 acetonitrile/triethylphosphate mobile phases.

Scheme I

styrene/divinylbenzene copolymer

- CH₂ - CH- CH₂

vinylpyridine/divinylbenzene copolymer

Both approaches were critically evaluated to see if the difference in the logarithm of the capacity factors ($\Delta \log k' \equiv$ "comparative retention") on the dual columns or dual mobile phases was a linear function solely of the hydrogen-bond acidity of the solute. The contribution to the measured $\Delta \log k'$ for each of the various factors that influence retention, such as solute size ($V_2/100$), dipolarity (π_2^*), polarizability (δ_2), hydrogen-bond acidity (α_2) and hydrogen-bond basicity (β_2), was investigated by chromatographing selected series of compounds. The influence of solute size on the

comparative retention was discerned by chromatographing a 4-alkylaniline homologous series. The effect of solute dipolarity and solute hydrogen-bond basicity was investigated by chromatographing a series of hydrogen-bond acceptor-only compounds that covered a wide range of dipolarity and hydrogen-bond basicity. The effect of solute polarizability was evaluated by chromatographing a series of polycyclic aromatic compounds. The dependence of the comparative retention on hydrogen-bond acidity was tested by chromatographing compounds ranging in acidity from aniline to p-nitrophenol.

For each test series of compounds, linear regression analysis was performed to determine the multiplicative coefficient for the parameter being probed. Multiple linear regression analysis was used to determine the coefficients for those series of compounds in which more than one factor was being probed and statistical tests on the regression variance were used to determine the statistical validity of the effect of each factor on $\Delta \log k'$. Values for the various solute parameters - $V_2/100$, π_2^* , δ_2 , α_2 , and β_2 - were taken from the literature.⁴ Since the solute parameter sets each cover the same range of values, direct comparison of the parameter coefficients allows for the assessment of the dependence of each factor on the comparative retention. As shown in Table I, for both approaches the solute hydrogen-bond acidity had the greatest effect on the comparative retention. However, the Δlog k' values determined from the dual mobile phase approach were less influenced by the solutes' nonhydrogen-bonding attributes and therefore, provide a purer measure of the solute hydrogen-bond acidity. Principle component analysis of the dual mobile phase $\Delta \log k'$ data for the hydrogen-bond donors indicates that over 99.9% of the variance is explained by one explanatory variable, which from the experimental results can be assigned to the solute hydrogen-bond acidity.

⁽a) Bondi, A. J. Phys. Chem. 1964, 68, 441. (b) Li, J.; Zhang, Y.; Dallas, A.J.; Carr, P.W. J. Chrom. 1991, 550, 101-134. (c) Kamlet, M.J.; Doherty, R.M.; Abraham, M.H.; Marcus, Y.; Taft, R.W. J. Phys. Chem. 1988, 92, 5244 - 5255. (d) Abraham, M.H.; Grellier, P.L.; Prior, D.V.; Duce, P.P.; Morris, J.J.; Taylor, P.J. J. Chem. Soc., Perkin Trans. 2 1989, 699-711. (e) Abraham, M.H.; Grellier, P.L.; Prior, D.V.; Morris, J.J.; Taylor, P.J. J. Chem. Soc., Perkin Trans. 2 1990, 521-529.

 $Table\ I$ Dependence of the Δ log k' Values on Various Solute Parameters

Solute	Matched Station	nary Phases Relative	Matched Mobile Phases Relative		
<u>Parameter</u>	Coefficient	<u>Value</u>	Coefficient	<u>Value</u>	
α 2	1.6	1.00	0.78	1.00	
β2	0.061	0.039	†	†	
π_2^*	0.19	0.12	-0.051	0.065	
δ_2	0.61	0.38	0.10	0.13	
V ₂ /100	-0.78	0.49	0.18	0.23	

[†] insignificant dependence on this variable as deemed by the Ehrenson F test.⁵

Currently, experiments are underway to utilize the dual mobile phase approach to assess the hydrogen-bond acidity of various hydrogen-bond donors used in cocrystallizations. Specific attention will be given to see if the expected hydrogen-bond associations based on the solute's measured hydrogen-bond acidity occur in the solid state.

⁵ Ehrenson, S. J. Org. Chem. 1979, 44, 1793-1797.

Professor O.T. Beachley, Jr. Department of Chemistry State University of New York Buffalo, NY 14214

Dr. Alan Berry Chemistry Division, Code 6120 Naval Research Laboratory 4555 Overlook Avenue, SW Washington, DC 20375-5000

Professor Jerald S. Bradshaw Department of Chemistry Brigham Young University Provo, UT 84602 R&T Code 413p002

Professor Ronald Breslow Department of Chemistry Columbia University New York, NY 10027 R&T Code 413p005

Dr. Duncan W. Brown Advanced Technology Materials, Inc. 520-B Danbury Road New Milford, CT 06776

Professor Herbert C. Brown Purdue University Department of Chemistry West Lafayette, IN 47907

Professor Steven L. Buchwald Department of Chemistry Massachusetts Institute of Tech Cambridge, MA 02139

Professor Cynthia J. Burrows
Department of Chemistry
State University of New York at
Stony Brook
Stony Brook, NY 11794-3400

Dr. Roque J. Calvo Executive Secretary The Electrochemical Society 10 South Main Street Pennington, NJ 08534-2896

Professor Peter Chen Department of Chemistry Harvard University Cambridge, MA 02138 Professor N. John Cooper Department of Chemistry University of Pittsburgh Pittsburgh, PA 15260

Professor Anthony W. Czarnik Department of Chemistry Ohio State University 120 West 18th Avenue Columbus, OH 43210-1173

Professor Peter Dervan Department of Chemistry California Institute of Technology Pasadena, CA 91125

Professor Francois N. Diederich Department of Chemistry University of California 405 Hilgard Avenue Los Angeles, CA 90024

Professor Dennis A. Dougherty Department of Chemistry California Institute of Technology Pasadena, CA 91125

Professor Kenneth M. Doxsee Department of Chemistry University of Oregon Eugene, OR 97403

Dr. Regis J. Ebner, Jr.
Director of Finance
Materials Research Society
9800 McKnight Road, Suite #27
Pittsburgh, PA 15237

Professor Margaret C. Etter Department of Chemistry University of Minnesota 207 Pleasant Street, S.E. Minneapolis, MN 55455

Professor Wilmer K. Fife Department of Chemistry Indiana University 1125 East 38th Street P.O. Box 647 Indianapolis, IN 46223

Professor Samuel H. Gellman Department of Chemistry University of Wisconsin Madison, WI 53706 Professor Andrew D. Hamilton Department of Chemistry University of Pittsburgh Pittsburgh, PA 15260

Prof. Mark J. Hampden-Smith Department of Chemistry University of New Mexico Albuquerque, NM 87131

Professor William E. Hatfield Department of Chemistry University of North Carolina Chapel Hill, NC 27514

Dr. Kelvin Higa Chemistry Division Research Department Naval Weapons Center China Lake, CA 93555

Professor Kenneth D. Karlin Merryman Hall 146 The Johns Hopkins University 34th & Charles Streets Baltimore, MD 21218

Professor Arthur E. Martell Department of Chemistry Texas A&M University College Station, TX 77843-3255

Professor Thomas J. McCarthy Department of Polymer Science University of Massachusetts Room 701 Graduate Research Center Amherst, MA 01003 R&T Code 400x015

Dr. Stephen W. McElvany Code 6113, Chemistry Division Naval Research Laboratory Washington, DC 20375-5000

Professor Lisa McElwee-White Department of Chemistry The Leland Stanford Junior Univ. Stanford, CA 94305

Professor Theordore G. Pavlopoulos Naval Ocean Systems Center Code 521 (B-111) San Diego, CA 92152-5000 Professor William S. Rees, Jr. Chemistry Division
The Florida State University
Tallahassee, FL 32306-3006

Professor Peter Schultz Department of Chemistry University of California Berkeley, CA 94720 R&T Code 413j005

Dr. Alok Singh Ctr. for Bio/Molec. Science Eng. Department of the Navy Naval Research Lab., Code 6090 Washingotn, DC 20375-5000

Dr. Michael L. Sinnott Unviersity of Brisol School of Chemistry Cantock's Close Briston, England BS8 1TS

Dr. Timothy M. Swager Departmentof Chemistry University of Pennsylvania Philadelphia, PA 19104-6323

Professor Richard L. Wells Department of Chemistry Duke University Durham, NC 27706

Professor Jeffrey D. Winkler Department of Chemistry The University of Chicago 5735 S. Ellis Avenue Chicago, IL 60637

Professor Jeffrey D. Zubkowski Department of Chemistry Jackson State University P.O. Box 17910, 1400 Lynch St. Jackson, MS 39217